

Gluon Radiation From Off-shell Quark in Glasma Fields

Based on: T. Lappi & IS arXiv:2607.XXXX


Ismail Soudi 

isma@physik.uni-bielefeld.de

Bielefeld University



Jets in heavy ion collisions



Medium

1. Initial Glasma Fields
2. Out-of-equilibrium evolution
2. Hydrodynamical QGP


Jet

1. Hard production $\tau_{\text{hard}} \sim 1/p_T$
2. Vacuum-like shower
3. Jet-medium Shower


HIC vs Small Systems

Heavy-ion collisions


- ▶ Strong signals of collectivity
- ▶ Clear quenching observed
- ▶ QGP energy loss dominates



PbPb



pPb



high-mult. pp

System size


- ▶ Energy loss v.s. system size
- ▶ Early-time v.s. late-time energy loss

[Chatrchyan & others Phys. Lett. B 718, 795–814 2013; Chatrchyan & others Phys. Lett. B 724, 213–240 2013; Khachatryan & others Phys. Lett. B 765, 193–220 2017; Hayrapetyan & others Phys. Rev. Lett. 136, 162301 2026]


HIC vs Small Systems

Heavy-ion collisions

- ▶ Strong signals of collective flow
- ▶ Clear quenching observed
- ▶ QGP energy loss dominates




PbPb



Small systems

- ▶ Like signals
- ▶ Quenching signal
- ▶ Correlations



high-mult. pp

▶ Energy loss v.s. system size

s. late-time energy loss

[Chatrchyan & others Phys. Lett. B 718, 795–814 2013; Chatrchyan & others Phys. Lett. B 724, 213–240 2013; Khachatryan & others Phys. Lett. B 765, 193–220 2017; Hayrapetyan & others Phys. Rev. Lett. 136, 162301 2026]

Momentum Broadening in Glasma






Figure 3: Schematic temporal evolution of $\hat{q}(t)$.

[Carrington et al. Phys. Lett. B 834, 137464 2022]



[Avramescu et al. Phys. Rev. D 107, 114021 2023]



[Ipp et al. Phys. Lett. B 810, 135810 2020]

► Large broadening $\hat{q}_{\text{glasma}} \gg \hat{q}_{\text{QGP}}$


► Short lifetime

► BDMPS-Z Energy loss: $\Delta E \sim \hat{q}L^2$

[Baier et al. Nucl. Phys. B 483, 291–320 1997]


Medium-Induced Radiation

Standard picture



- ▶ Medium length $L \gg 1/Q_s$
- ▶ Multiple scatterings (LPM)
- ▶ Vacuum/medium separation
→ Jet resolved as on-shell

Early-time CGC




- ▶ Short-lived glasma fields
- ▶ Highly virtual jet
- ▶ Vacuum-like splittings
→ Virtuality vs glasma resolution

First emission: vacuum-like vs glasma modification

$$\tau_f^{\text{vac}} \Leftrightarrow 1/Q_s.$$

Shockwave Framework

- ▶ Dense small- x gluons
 - Classical color field A^μ .
- ▶ Fast target is a thin sheet in x^+ .
 - $A^\mu(x) = \delta^{\mu-} \delta(x^+) \alpha(x)$.
- ▶ Crossing the sheet described by
 - Wilson line $\mathcal{U}(x)$.



- ▶ Frozen background field during the short interaction.

[Balitsky Nucl. Phys. B 463, 99–160 1996; Kovchegov Phys. Rev. D 61, 74018 2000; Iancu & Venugopalan Quark-gluon plasma 4 249–3363 2003; Gelis & Jalilian-Marian Phys. Rev. D 66, 14021 2002]

LCPT

- ▶ In light-cone gauge $A^+ = 0$

$$\begin{aligned} S_F(x, y) &= \int dq^- e^{i(x^+ - y^+) \cdot q^-} \frac{\not{q}}{q^2 + i\varepsilon} \\ &= \int dq^- e^{i(x^+ - y^+) \cdot q^-} \frac{1}{2q^+} \frac{\not{q}}{q^- - \frac{\mathbf{q}}{2q^+} + i\varepsilon} + \frac{\gamma^+ \left(q^- - \frac{\mathbf{q}}{2q^+} \right)}{2q^+ \left(q^- - \frac{\mathbf{q}}{2q^+} \right)} \\ &= \theta(x^+ - y^+) \frac{1}{2q^+} \frac{\not{q}}{q^- - \frac{\mathbf{q}}{2q^+} + i\varepsilon} + \frac{\gamma^+}{2q^+} \delta(x^+ - y^+) + \text{Reverse Order} \end{aligned} \tag{1}$$

- ▶ Old-fashioned perturbation theory with x^+ time-ordering (c.f. *T. Lappi's* talk)

Quark Propagator

Eikonal propagation

$$S_{F(x,y)_{\beta\alpha}} = \text{Diagram: A quark line from } x \text{ to } y \text{ with a vertical red shaded region in the middle. The line has arrows pointing right, labeled } p \text{ and } k. \text{ The shaded region contains a circle with a cross.}$$

$$= \mathbf{1}_{\beta\alpha} \delta^{(2)}(\mathbf{x} - \mathbf{y}) \delta(x^+ - y^+) \text{sign}(x^- - y^-) \frac{\gamma^+}{4} + \int \frac{d^3 p}{(2\pi)^3} \int \frac{d^3 k}{(2\pi)^3} \quad (2)$$

$$\frac{\bar{p} \gamma^+ \bar{k}}{4p^+ k^+} (2\pi) \delta(p^+ - k^+) e^{-ix \cdot \bar{p} + iy \cdot \bar{k}} \int d^2 z e^{-iz \cdot (p - k)} \\ \times \left\{ \theta(k^+) \theta(x^+ - y^+) \mathcal{U}_F(x^+, y^+; z)_{\beta\alpha} - \theta(-k^+) \theta(y^+ - x^+) \mathcal{U}_F(y^+, x^+; z)_{\beta\alpha} \right\}.$$

[Balitsky Nucl. Phys. B 463, 99–160 1996; Kovchegov Phys. Rev. D 61, 74018 2000; Iancu & Venugopalan Quark-gluon plasma 4 249–3363 2003; Gelis & Jalilian-Marian Phys. Rev. D 66, 14021 2002; Gelis & Venugopalan Phys. Rev. D 69, 14019 2004; Fujii et al. Nucl. Phys. A 780, 146–174 2006]

Gluon Propagator

Adjoint representation

$$\begin{aligned}
 G_{ab}^{\mu\nu}(x, y) &= \text{Diagram: A wavy line with momentum } p \text{ from point } x \text{ to a red shaded box containing a crossed circle, followed by a wavy line with momentum } k \text{ to point } y. \\
 &= G_{ab, \text{inst}}^{\mu\nu}(x, y) + \int \frac{d^3 p}{(2\pi)^3} \int \frac{d^3 k}{(2\pi)^3} \frac{1}{2k^+} \sum_{\lambda} \varepsilon_{\lambda}^{\mu}(\bar{p}) \varepsilon_{\lambda}^{*\nu}(\bar{k}) \\
 &\quad \times (2\pi) \delta(p^+ - k^+) e^{-ix \cdot \bar{p} + iy \cdot \bar{k}} \int d^2 z e^{-iz \cdot (p - k)} \\
 &\quad \times \left\{ \theta(k^+) \theta(x^+ - y^+) \mathcal{U}_A(x^+, y^+; z)_{ab} - \theta(-k^+) \theta(y^+ - x^+) \mathcal{U}_A(y^+, x^+; z)_{ab} \right\}.
 \end{aligned} \tag{3}$$

[Balitsky Nucl. Phys. B 463, 99–160 1996; Kovchegov Phys. Rev. D 61, 74018 2000; Iancu & Venugopalan Quark-gluon plasma 4 249–3363 2003; Gelis & Jalilian-Marian Phys. Rev. D 66, 14021 2002; Gelis & Venugopalan Phys. Rev. D 69, 14019 2004; Fujii et al. Nucl. Phys. A 780, 146–174 2006]

Kinematics

- Incoming hard quark:


$$p_1 = (0^+, p_1^-, \mathbf{0}), \quad \ell = (\ell^+, \ell^-, \mathbf{0}) \quad (4)$$

- On-shell Outgoing pair:

$$p' = (p'^+, p'^-, \mathbf{p}'), \quad k' = (k'^+, k'^-, \mathbf{k}') \quad (5)$$

- Gluon momentum fraction

$$y = \frac{k'^+}{q'^+} = \frac{k'^+}{p'^+ + k'^+} \quad (6)$$





b : quark position

r : gluon position

$u = r - b$: qg size

On-Shell Case



► **A**: Emission **after** the shockwave

► **B**: Emission **before** the shockwave

► On-shell parent $\rightarrow$ two time orderings across the shockwave.

The Wave Function

- ▶ The $q \rightarrow qg$ wave function

$$\Psi^{q \rightarrow qg}(p', k') = \begin{array}{c} \text{Diagram: A horizontal line with a black dot at position } z_1. \text{ An incoming arrow from the left is labeled } q'. \text{ An outgoing arrow to the right is labeled } p'. \text{ A wavy line (gluon) goes from the dot to the upper right, labeled } k'. \end{array} = \frac{\bar{u}(q' - k') \not{\epsilon}_\lambda^*(k') u(q')}{(q' - k')^- - q'^- - k'^-} \quad (7)$$


- ▶ On-shell parent $\rightarrow$ Energy denominator simply: $(k - yq')^{-2} \neq \infty$.
- ▶ Fourier Transform $q' \rightarrow r - b$:

$$\begin{aligned} \phi_0^\lambda(u) &= \int d^2 q' e^{i u \cdot q'} \Psi^{q \rightarrow qg}(p', k') \\ &= 2i \sqrt{1-y} \frac{u \cdot \epsilon_\lambda^*}{2\pi u^2} [\delta_{\lambda,s} + (1-y)\delta_{\lambda,-s}]. \end{aligned} \quad (8)$$

[Marquet Nucl. Phys. A 796, 41–60 2007]

Amplitude

After




$$\mathcal{A}_{\text{after}} = \quad (9)$$

$$= -\mathcal{N}_0 \int d^2b d^2r e^{-i\mathbf{b} \cdot \mathbf{p}'} e^{-i\mathbf{r} \cdot \mathbf{k}'} \phi_0^\lambda(\mathbf{r} - \mathbf{b}) t^a \mathcal{U}_F(\mathbf{v}).$$

► Transverse center of mass

→ $\mathbf{v} = \mathbf{b} + y(\mathbf{r} - \mathbf{b}).$

Before



$$\mathcal{A}_{\text{before}} = \quad (10)$$

$$= \mathcal{N}_0 \int d^2b d^2r e^{-i\mathbf{b} \cdot \mathbf{p}'} e^{-i\mathbf{r} \cdot \mathbf{k}'} \phi_0^\lambda(\mathbf{r} - \mathbf{b}) t^b \mathcal{U}_F(\mathbf{b}) \mathcal{U}_A^{ba}(\mathbf{r}).$$

► Transverse positions:

→ $\mathbf{b}$: quark

→ $\mathbf{r}$: gluon

Total Amplitude

$$\mathcal{A}_{\text{on-shell}} = \mathcal{N}_0 \int d^2b d^2r e^{-i\delta\mathbf{r} \cdot (\mathbf{p}' + \mathbf{k}')} e^{-i\mathbf{R} \cdot \frac{\mathbf{k}' - \mathbf{p}'}{2}} \phi_0^\lambda(\delta\mathbf{r})$$

$$\left[\underbrace{t^b \mathcal{U}_F(\mathbf{b}) \mathcal{U}_A(\mathbf{r})^{ba}}_{\text{After}} - \underbrace{t^a \mathcal{U}_F(\mathbf{v})}_{\text{Before}} \right]. \quad (11)$$

- ▶ Amplitude $\rightarrow$ difference between **Before** and **After**.
- ▶ In vacuum: $\rightarrow \mathcal{U} \rightarrow 1 \Rightarrow \mathcal{A} \rightarrow 0.$

Kinematics

- Incoming hard quark:


$$p_1 = (0^+, p_1^-, \mathbf{0}), \quad \ell = (\ell^+, \ell^-, \mathbf{0}) \quad (12)$$

- On-shell Outgoing pair:

$$p' = (p'^+, p'^-, \mathbf{p}'), \quad k' = (k'^+, k'^-, \mathbf{k}') \quad (13)$$

- Gluon momentum fraction

$$y = \frac{k'^+}{q'^+} = \frac{k'^+}{p'^+ + k'^+} \quad (14)$$



b : quark position


r : gluon position

$u = r - b$: qg size

Off-shell Amplitude


- **A:** emission **after** the shockwave

$$y_1^+ < 0 < z_1^+$$




- **B:** emission **before** the shockwave

$$y_1^+ < z_1^+ < 0$$



- **C:** shockwave **before** the photon vertex


$$0 < y_1^+ < z_1^+$$



Vacuum radiation

► Minus energy

• $z_1^+ : P^- = \bar{p}^- + \bar{k}^-$ • $y_1^+ : Q^- = p_1^- + \ell^-$



► **A:** $y_1^+ < 0 < z_1^+$

$$\int_{-\infty}^0 dy_1^+ \int_0^{\infty} dz_1^+ \mathcal{V}$$

$$= - \frac{1}{P^- - \bar{q}^- + i\varepsilon} \frac{1}{Q^- - \bar{q}^- + i\varepsilon}$$

► **B:** $y_1^+ < z_1^+ < 0$

$$\int_{-\infty}^0 dy_1^+ \int_{y_1^+}^0 dz_1^+ \mathcal{V}$$

$$= \frac{1}{P^- - Q^- - i\varepsilon} \frac{1}{Q^- - \bar{q}^- + i\varepsilon}$$

► **C:** $0 < y_1^+ < z_1^+$

$$\int_0^{\infty} dy_1^+ \int_{y_1^+}^{\infty} dz_1^+ \mathcal{V}$$

$$= \frac{1}{P^- - \bar{q}^- + i\varepsilon} \frac{1}{Q^- - P^- - i\varepsilon}$$

Energy conservation

- The sum of the three time-orderings gives a delta function in minus energy

$$\begin{aligned}
 & \frac{1}{P^- - \bar{q}^- + i\varepsilon} \frac{1}{Q^- - P^- - i\varepsilon} - \frac{1}{P^- - \bar{q}^- + i\varepsilon} \frac{1}{Q^- - \bar{q}^- + i\varepsilon} \\
 & + \frac{1}{P^- - Q^- - i\varepsilon} \frac{1}{Q^- - \bar{q}^- + i\varepsilon} \\
 & = \frac{1}{Q^- - \bar{q}^- + i\varepsilon} \left[\frac{1}{Q^- - P^- - i\varepsilon} - \frac{1}{Q^- - P^- + i\varepsilon} \right] \propto \frac{1}{Q^- - \bar{q}^- + i\varepsilon} \delta(Q^- - P^-)
 \end{aligned} \tag{15}$$

Emission **After** Shockwave

- ▶ Photon vertex:


$$\bar{u}(\bar{q}) \not{\epsilon}_\lambda(\ell) u(\bar{p}) = 2\sqrt{p_1^-} \ell^+$$

- ▶ Energy denominator ($q = 0$):

$$p_1^- + \ell^- - \bar{p}^- - \bar{k}^- + i\varepsilon = -\frac{k^2}{2\ell^+ y(1-y)} \quad (17)$$

- ▶ Weizsäcker-Williams wave function:

$$\phi_0^\lambda(\mathbf{u}) = 2i\sqrt{1-y} \frac{\mathbf{u} \cdot \boldsymbol{\varepsilon}_\lambda^*}{2\pi u^2} [\delta_{\lambda,s} + (1-y)\delta_{\lambda,-s}] \quad (18)$$




Emission **After** Shockwave

- Weizsäcker-Williams wave function:

$$\phi_0^\lambda(\mathbf{u}) = 2i\sqrt{1-y} \frac{\mathbf{u} \cdot \boldsymbol{\varepsilon}_\lambda^*}{2\pi \mathbf{u}^2} [\delta_{\lambda,s} + (1-y)\delta_{\lambda,-s}] \quad (19)$$

- Amplitude for ordering A:

$$\mathcal{A}_A = -2eg \frac{\sqrt{p_1^- \ell^+}}{p_1^- + \ell^-} \int d^2b d^2r e^{-i\mathbf{b} \cdot \mathbf{p}'} e^{-i\mathbf{r} \cdot \mathbf{k}'} \phi_0^\lambda(\mathbf{u}) t^a \mathcal{U}_F(\mathbf{v})_{\beta\alpha} . \quad (20)$$



Emission **Before** Shockwave

- Off-shell virtuality:


$$M^2 = y(1 - y)(2\ell^+ p_1^- - Q^2)$$

- Energy denominator ($q = 0$):

$$p_1^- + \ell^- - \bar{p}^- - \bar{k}^- + i\varepsilon = -\frac{k^2 - M^2 - i\varepsilon}{2\ell^+ y(1 - y)} \quad (21)$$

- $K_1(\sqrt{-M^2 - i\varepsilon}) \rightarrow i\varepsilon$ selects the side of **Branch cut**

$$K_1(ix) = \begin{cases} H^{(1)}(x) = -\frac{\pi}{2}[J_1(x) + iY_1(x)], & -\pi < \arg(x) < -\frac{\pi}{2} \\ H^{(2)}(x) = -\frac{\pi}{2}[J_1(x) - iY_1(x)], & -\frac{\pi}{2} < \arg(x) < \pi \end{cases} \quad (22)$$



Emission **Before** Shockwave

- $K_1\left(\sqrt{-M^2 - i\varepsilon}\right) \rightarrow i\varepsilon$ selects the side of **Branch cut**


$$K_1(ix) = \begin{cases} H^{(1)}(x) = -\frac{\pi}{2}[J_1(x) + iY_1(x)], & -\pi < \arg(x) < -\frac{\pi}{2} \\ H^{(2)}(x) = -\frac{\pi}{2}[J_1(x) - iY_1(x)], & -\frac{\pi}{2} < \arg(x) < \pi \end{cases} \quad (23)$$

- wave function:

$$\phi_-^\lambda(\mathbf{u}) = i \frac{(-iM)H^{(2)}(-iM|\mathbf{u}|)}{2\pi} \frac{\mathbf{u} \cdot \boldsymbol{\varepsilon}_\lambda^*}{|\mathbf{u}|} 2\sqrt{1-y} [\delta_{\lambda,s} + (1-y)\delta_{\lambda,-s}] \quad (24)$$

- Amplitude:

$$\mathcal{A}_B = 2eg \frac{\sqrt{p_1^- \ell^+}}{p_1^- + \ell^-} \int d^2b d^2r e^{-i\mathbf{b} \cdot \mathbf{p}'} e^{-i\mathbf{r} \cdot \mathbf{k}'} \phi_-^\lambda(\mathbf{u}) t^b \mathcal{U}_F(\mathbf{b})_{\beta\alpha} \mathcal{U}_A^{ba}(\mathbf{r}). \quad (25)$$



Shockwave **Before** Photon Vertex

- ▶ Off-shell virtuality:


$$\sqrt{-M^2 + i\varepsilon} = iM$$

- ▶ Energy denominator prescription:

$$p_1^- + \ell^- - p'^- - k'^- - i\varepsilon = -\frac{\mathbf{k}'^2 - M^2 + i\varepsilon}{2\ell^+ y(1-y)} \quad (27)$$

- ▶ Amplitude contains both ϕ_+ and vacuum ϕ_0 :


$$\frac{1}{p_1^- + \ell^- - p'^- - k'^- - i\varepsilon} \frac{\mathcal{N}}{2\ell^+} = - \int d^2\mathbf{r} e^{-i\mathbf{r} \cdot \mathbf{k}'} [\phi_+^\lambda(\mathbf{u}) - \phi_0^\lambda(\mathbf{u})] \quad (28)$$



Shockwave **Before** Photon Vertex

- ▶ Off-shell virtuality:

$$\sqrt{-M^2 + i\varepsilon} = iM$$



- ▶ Energy denominator prescription:

$$p_1^- + \ell^- - p'^- - k'^- - i\varepsilon = -\frac{k'^2 - M^2 + i\varepsilon}{2\ell^+ y(1 - y)} \quad (30)$$

- ▶ Amplitude for ordering C:

$$\mathcal{A}_C = 2eg \frac{\sqrt{p_1^- \ell^+}}{p_1^- + \ell^-} \int d^2 b d^2 r e^{-i\mathbf{b} \cdot \mathbf{p}'} e^{-i\mathbf{r} \cdot \mathbf{k}'} [\phi_0^\lambda(\mathbf{u}) - \phi_+^\lambda(\mathbf{u})] t^a \mathbf{1}_{\beta\alpha}. \quad (31)$$

Total Amplitude




- Use $\mathbf{u} = \mathbf{r} - \mathbf{b}$ and $\mathbf{v} = \mathbf{b} + y(\mathbf{r} - \mathbf{b})$:

$$\mathbf{b} = \mathbf{v} - y\mathbf{u}, \quad \mathbf{r} = \mathbf{v} + (1 - y)\mathbf{u}, \quad (32)$$

- Total amplitude:

$$\mathcal{A}_{\text{total}} = 2eg \frac{\sqrt{p_1^- \ell^+}}{p_1^- + \ell^-} \int d^2\mathbf{u} d^2\mathbf{v} e^{-i\mathbf{v} \cdot (\mathbf{p}' + \mathbf{k}')} e^{-i\mathbf{u} \cdot ((1-y)\mathbf{k}' - y\mathbf{p}')} \quad (33)$$

$$\times \left[\underbrace{\phi_-^\lambda(\mathbf{u}) t^b \mathcal{U}_F(\mathbf{b})_{\beta\alpha} \mathcal{U}_A(\mathbf{r})^{ba}}_{\mathbf{B}} - \underbrace{\phi_0^\lambda(\mathbf{u}) t^a \mathcal{U}_F(\mathbf{v})_{\beta\alpha}}_{\mathbf{A}} + \underbrace{[\phi_0^\lambda(\mathbf{u}) - \phi_+^\lambda(\mathbf{u})] t^a \mathbf{1}_{\beta\alpha}}_{\mathbf{C}} \right]$$



Vacuum Limit

- ▶ Taking $\mathcal{U}_{F/A} \rightarrow 1$:

$$\mathcal{A}_{\text{total}} = 2eg \frac{\sqrt{p_1^- \ell^+}}{p_1^- + \ell^-} \int d^2 \mathbf{u} d^2 \mathbf{v} e^{-i\mathbf{v} \cdot (\mathbf{p}' + \mathbf{k}')} e^{-i\mathbf{u} \cdot ((1-y)\mathbf{k}' - y\mathbf{p}')} t^a [\phi_-^\lambda(\mathbf{u}) - \phi_+^\lambda(\mathbf{u})]. \quad (34)$$

→ On-shell contribution $\nrightarrow \phi_0 \rightarrow 0$.

- ▶ Integral $\int d\mathbf{v} \rightarrow (2\pi)^2 \delta^{(2)}(\mathbf{p}' + \mathbf{k}')$, transverse momentum conservation.
- ▶ Difference $\phi_-^\lambda - \phi_+^\lambda \rightarrow$ minus-momentum conservation

$$\phi_-^\lambda - \phi_+^\lambda = -\phi_J^\lambda(\mathbf{u})$$

$$\text{Momentum space} \Rightarrow \frac{1}{\mathbf{k}^2 - M^2 - i\varepsilon} - \frac{1}{\mathbf{k}^2 - M^2 + i\varepsilon} = 2\pi i \delta(\mathbf{k}^2 - M^2) \quad (35)$$

Vacuum Subtraction

- ▶ The vacuum square contains the usual square of an S -matrix delta function:

$$\int d^2u d^2\bar{u} e^{-il \cdot u} e^{il \cdot \bar{u}} \phi_J^\lambda(u) \phi_J^{\lambda*}(\bar{u}) \propto [\delta(l^2 - M^2)]^2 \quad (36)$$

- ▶ Organize the finite medium-dependent object as

$$\langle |\mathcal{A}|^2 \rangle - |\mathcal{A}_{\text{vac}}|^2 = \langle |\mathcal{A}_{\text{med}}|^2 \rangle + 2 \text{Re} \langle \mathcal{A}_{\text{vac}} \mathcal{A}_{\text{med}}^\dagger \rangle, \quad \mathcal{A}_{\text{med}} = \mathcal{A} - \mathcal{A}_{\text{vac}}. \quad (37)$$

Medium Average

- Dipole and multipole correlators:

$$\begin{aligned} S_{q\bar{q}}^2(\mathbf{v}, \bar{\mathbf{v}}) &= \frac{1}{N_c} \langle \text{Tr} [\mathcal{U}_{F(\mathbf{v})} \mathcal{U}_F^{\dagger(\bar{\mathbf{v}})}] \rangle \\ S_{qg\bar{q}}^3(\mathbf{b}, \mathbf{r}, \bar{\mathbf{v}}) &= \frac{1}{C_F N_c} \langle \text{Tr} [t^b \mathcal{U}_{F(\mathbf{b})} t^a \mathcal{U}_F^{\dagger(\bar{\mathbf{v}})}] \mathcal{U}_{A(\mathbf{r})}^{ba} \rangle \\ S_{qg\bar{q}g}^4(\mathbf{b}, \bar{\mathbf{b}}, \mathbf{r}, \bar{\mathbf{r}}) &= \frac{1}{C_F N_c} \langle \text{Tr} [\mathcal{U}_{F(\mathbf{b})} \mathcal{U}_F^{\dagger(\bar{\mathbf{b}})} t^a t^b] \mathcal{U}_{A(\mathbf{r})}^{ac} \mathcal{U}_A^{\dagger(\bar{\mathbf{r}})cb} \rangle \end{aligned} \tag{38}$$

Square of the Amplitude

$$\begin{aligned} \langle |\mathcal{A}_{\text{med}}|^2 \rangle &= 4e^2 g^2 C_F N_c \frac{p_1^- \ell^+}{(p_1^- + \ell^-)^2} \int d^2 \boldsymbol{v} d^2 \bar{\boldsymbol{v}} d^2 \boldsymbol{u} d^2 \bar{\boldsymbol{u}} e^{-i(\boldsymbol{v} - \bar{\boldsymbol{v}}) \cdot (\boldsymbol{p}' + \boldsymbol{k}')} e^{-i(\boldsymbol{u} - \bar{\boldsymbol{u}}) \cdot ((1-y)\boldsymbol{k}' - y\boldsymbol{p}')} \\ &\times \left\{ \phi_0^\lambda(\boldsymbol{u}) \phi_0^{\lambda*}(\bar{\boldsymbol{u}}) [S^4(\boldsymbol{b}, \bar{\boldsymbol{b}}, \boldsymbol{r}, \bar{\boldsymbol{r}}) - S^3(\boldsymbol{b}, \boldsymbol{r}, \bar{\boldsymbol{v}}) - S^3(\bar{\boldsymbol{b}}, \bar{\boldsymbol{r}}, \boldsymbol{v}) + S^2(\boldsymbol{v}, \bar{\boldsymbol{v}})] \right\} \end{aligned} \quad (39)$$

Square of the Amplitude

$$\begin{aligned}
 \langle |\mathcal{A}_{\text{med}}|^2 \rangle &= 4e^2 g^2 C_F N_c \frac{p_1^- \ell^+}{(p_1^- + \ell^-)^2} \int d^2 v d^2 \bar{v} d^2 u d^2 \bar{u} e^{-i(v-\bar{v}) \cdot (p' + k')} e^{-i(u-\bar{u}) \cdot ((1-y)k' - y p')} \\
 &\times \left\{ \phi_0^\lambda(u) \phi_0^{\lambda*}(\bar{u}) [S^4(b, \bar{b}, r, \bar{r}) - S^3(b, r, \bar{v}) - S^3(\bar{b}, \bar{r}, v) + S^2(v, \bar{v})] \right. \\
 &\quad + [\phi_-^\lambda(u) - \phi_0^\lambda(u)] \phi_0^{\lambda*}(\bar{u}) [S^4(b, \bar{b}, r, \bar{r}) - S^3(b, r, \bar{v})] \\
 &\quad + \phi_0^\lambda(u) [\phi_-^{\lambda*}(\bar{u}) - \phi_0^{\lambda*}(\bar{u})] [S^4(b, \bar{b}, r, \bar{r}) - S^3(\bar{b}, \bar{r}, v)] \\
 &\quad \left. + [\phi_-^\lambda(u) - \phi_0^\lambda(u)] [\phi_-^{\lambda*}(\bar{u}) - \phi_0^{\lambda*}(\bar{u})] [S^4(b, \bar{b}, r, \bar{r}) + 1] \right\}.
 \end{aligned} \tag{40}$$

Square of the Amplitude

$$\begin{aligned}
 \langle |\mathcal{A}_{\text{med}}|^2 \rangle &= 4e^2 g^2 C_F N_c \frac{p_1^- \ell^+}{(p_1^- + \ell^-)^2} \int d^2 v d^2 \bar{v} d^2 u d^2 \bar{u} e^{-i(v-\bar{v}) \cdot (p' + k')} e^{-i(u-\bar{u}) \cdot ((1-y)k' - y p')} \\
 &\times \left\{ \phi_0^\lambda(u) \phi_0^{\lambda*}(\bar{u}) [S^4(b, \bar{b}, r, \bar{r}) - S^3(b, r, \bar{v}) - S^3(\bar{b}, \bar{r}, v) + S^2(v, \bar{v})] \right. \\
 &\quad + [\phi_-^\lambda(u) - \phi_0^\lambda(u)] \phi_0^{\lambda*}(\bar{u}) [S^4(b, \bar{b}, r, \bar{r}) - S^3(b, r, \bar{v})] \\
 &\quad + \phi_0^\lambda(u) [\phi_-^{\lambda*}(\bar{u}) - \phi_0^{\lambda*}(\bar{u})] [S^4(b, \bar{b}, r, \bar{r}) - S^3(\bar{b}, \bar{r}, v)] \\
 &\quad \left. + [\phi_-^\lambda(u) - \phi_0^\lambda(u)] [\phi_-^{\lambda*}(\bar{u}) - \phi_0^{\lambda*}(\bar{u})] [S^4(b, \bar{b}, r, \bar{r}) + 1] \right\}. \tag{41}
 \end{aligned}$$

► Setting $M \rightarrow 0$ gives $[\phi_-^\lambda(u) - \phi_0^\lambda(u)] \rightarrow 0$

→ recovering [Marquet Nucl. Phys. A 796, 41–60 2007]

Relevant Scales

- ▶ The wave function

$$\phi_-^\lambda(\boldsymbol{u}) \propto M[J_1(Mu) - iY_1(Mu)] \frac{\boldsymbol{u} \cdot \boldsymbol{\varepsilon}_\lambda^*}{u}, \quad (42)$$

→ Selects transverse size $u \approx 1/M$.

- ▶ Dipole correlator

$$S(u) \approx \exp\left(-Q_s^2 \frac{u^2}{4}\right). \quad (43)$$

→ Resolves transverse sizes $u \approx 1/Q_s$.

Small Virtuality $M^2 \ll Q_s^2$

- ▶ Dipole correlators sets transverse size

$$u \approx 1/Q_s \quad \Rightarrow \quad Mu \ll 1. \quad (44)$$

- ▶ Small- Mu expansion:

$$J_1(M|u|) \approx 0, \quad Y_1(M|u|) \approx -\frac{2}{\pi M|u|}, \quad \Rightarrow \quad \phi_-^\lambda(\mathbf{u}) = \phi_0^\lambda(\mathbf{u}). \quad (45)$$

The square matrix element reduces to the on-shell case.

$$\phi_0^\lambda(u) \phi_0^{\lambda*}(\bar{u}) [S^4(b, \bar{b}, r, \bar{r}) - S^3(b, r, \bar{v}) - S^3(\bar{b}, \bar{r}, v) + S^2(v, \bar{v})] \quad (46)$$

High Virtuality $M^2 \gg Q_s^2$

- ▶ Large- Mu Hankel phase:

$$H_1^{(2)}(M|u|) \sim \frac{e^{-iM|u|}}{\sqrt{M|u|}}, \quad M \gg Q_s \Rightarrow u \approx 1/M \ll 1/Q_s. \quad (47)$$

- ▶ Small dipole:

$$S(u) = \exp\left(-Q_s^2 \frac{u^2}{4}\right) \approx 1 - Q_s^2 \frac{u^2}{4}, \quad 1 - S(u) \approx Q_s^2/(4M^2). \quad (48)$$

- ▶ Medium correction:

$$\frac{|\mathcal{A}_{\text{med}}|^2}{|\mathcal{A}_{\text{vac}}|^2} \approx \mathcal{O}\left(\frac{Q_s^2}{M^2}\right). \quad (49)$$

- ▶ Vacuum-like radiation dominates;
→ Suppressed medium corrections.

Intermediate Virtuality $M^2 \approx Q_s^2$

- ▶ Two transverse scales

$$u \approx 1/Q_s \approx 1/M, \quad Mu \approx 1. \quad (50)$$

- ▶ Important contributions from all terms

$$\begin{aligned} & \phi_0^\lambda(u) \phi_0^{\lambda*}(\bar{u}) [S^4(b, \bar{b}, r, \bar{r}) - S^3(b, r, \bar{v}) - S^3(\bar{b}, \bar{r}, v) + S^2(v, \bar{v})] \\ & + [\phi_-^\lambda(u) - \phi_0^\lambda(u)] \phi_0^{\lambda*}(\bar{u}) [S^4(b, \bar{b}, r, \bar{r}) - S^3(b, r, \bar{v})] \\ & + \phi_0^\lambda(u) [\phi_-^{\lambda*}(\bar{u}) - \phi_0^{\lambda*}(\bar{u})] [S^4(b, \bar{b}, r, \bar{r}) - S^3(\bar{b}, \bar{r}, v)] \\ & + [\phi_-^\lambda(u) - \phi_0^\lambda(u)] [\phi_-^{\lambda*}(\bar{u}) - \phi_0^{\lambda*}(\bar{u})] [S^4(b, \bar{b}, r, \bar{r}) + 1]. \end{aligned} \quad (51)$$

- ▶ Interplay between Glasma scales and virtuality.

Conclusions

Takeaways

- ▶ Off-shell radiation introduces a new hard scale M^2 in the shockwave calculation.
- ▶ The glasma resolves transverse sizes $u \approx 1/Q_s$:

$$M^2 \ll Q_s^2 \Rightarrow \text{on-shell limit}, \quad M^2 \gg Q_s^2 \Rightarrow \text{color transparency}. \quad (52)$$

- ▶ The glasma selects part of early shower with virtuality of order $\sim Q_s^2$.

Outlook

- ▶ Compute the medium-induced spectrum $\frac{dN}{dy d^2\mathbf{k}'}$ and compare to on-shell results.
- ▶ Understand interference term $2 \operatorname{Re} \langle \mathcal{A}_{\text{vac}} \mathcal{A}_{\text{med}}^\dagger \rangle$.

Bibliography

- [1] S. Chatrchyan and others, “Observation of Long-Range Near-Side Angular Correlations in Proton-Lead Collisions at the LHC,” *Phys. Lett. B*, vol. 718, pp. 795–814, 2013, doi: [10.1016/j.physletb.2012.11.025](https://doi.org/10.1016/j.physletb.2012.11.025).
- [2] S. Chatrchyan and others, “Multiplicity and Transverse Momentum Dependence of Two- and Four-Particle Correlations in pPb and PbPb Collisions,” *Phys. Lett. B*, vol. 724, pp. 213–240, 2013, doi: [10.1016/j.physletb.2013.06.028](https://doi.org/10.1016/j.physletb.2013.06.028).
- [3] V. Khachatryan and others, “Evidence for collectivity in pp collisions at the LHC,” *Phys. Lett. B*, vol. 765, pp. 193–220, 2017, doi: [10.1016/j.physletb.2016.12.009](https://doi.org/10.1016/j.physletb.2016.12.009).
- [4] A. Hayrapetyan and others, “Observation of Suppressed Charged-Particle Production in Ultrarelativistic Oxygen-Oxygen Collisions,” *Phys. Rev. Lett.*, vol. 136, no. 16, p. 162301, 2026, doi: [10.1103/89sf-9t1x](https://doi.org/10.1103/89sf-9t1x).
- [5] M. E. Carrington, A. Czajka, and S. Mrowczynski, “Jet quenching in glasma,” *Phys. Lett. B*, vol. 834, p. 137464, 2022, doi: [10.1016/j.physletb.2022.137464](https://doi.org/10.1016/j.physletb.2022.137464).
- [6] D. Avramescu, V. Băran, V. Greco, A. Ipp, D. I. Müller, and M. Ruggieri, “Simulating jets and heavy quarks in the glasma using the colored particle-in-cell method,” *Phys. Rev. D*, vol. 107, no. 11, p. 114021, 2023, doi: [10.1103/PhysRevD.107.114021](https://doi.org/10.1103/PhysRevD.107.114021).
- [7] A. Ipp, D. I. Müller, and D. Schuh, “Jet momentum broadening in the pre-equilibrium Glasma,” *Phys. Lett. B*, vol. 810, p. 135810, 2020, doi: [10.1016/j.physletb.2020.135810](https://doi.org/10.1016/j.physletb.2020.135810).
- [8] R. Baier, Y. L. Dokshitzer, A. H. Mueller, S. Peigne, and D. Schiff, “Radiative energy loss of high-energy quarks and gluons in a finite volume quark - gluon plasma,” *Nucl. Phys. B*, vol. 483, pp. 291–320, 1997, doi: [10.1016/S0550-3213\(96\)00553-6](https://doi.org/10.1016/S0550-3213(96)00553-6).
- [9] I. Balitsky, “Operator expansion for high-energy scattering,” *Nucl. Phys. B*, vol. 463, pp. 99–160, 1996, doi: [10.1016/0550-3213\(95\)00638-9](https://doi.org/10.1016/0550-3213(95)00638-9).
- [10] Y. V. Kovchegov, “Unitarization of the BFKL pomeron on a nucleus,” *Phys. Rev. D*, vol. 61, p. 74018, 2000, doi: [10.1103/PhysRevD.61.074018](https://doi.org/10.1103/PhysRevD.61.074018).
- [11] E. Iancu and R. Venugopalan, “The Color glass condensate and high-energy scattering in QCD,” in *Quark-gluon plasma 4*, R. C. Hwa and X.-N. Wang, Eds., 2003, pp. 249–3363. doi: [10.1142/9789812795533_0005](https://doi.org/10.1142/9789812795533_0005).
- [12] F. Gelis and J. Jalilian-Marian, “Photon production in high-energy proton nucleus collisions,” *Phys. Rev. D*, vol. 66, p. 14021, 2002, doi: [10.1103/PhysRevD.66.014021](https://doi.org/10.1103/PhysRevD.66.014021).

Bibliography

- [13] F. Gelis and R. Venugopalan, “Large mass q anti- q production from the color glass condensate,” *Phys. Rev. D*, vol. 69, p. 14019, 2004, doi: [10.1103/PhysRevD.69.014019](https://doi.org/10.1103/PhysRevD.69.014019).
- [14] H. Fujii, F. Gelis, and R. Venugopalan, “Quark pair production in high energy pA collisions: General features,” *Nucl. Phys. A*, vol. 780, pp. 146–174, 2006, doi: [10.1016/j.nuclphysa.2006.09.012](https://doi.org/10.1016/j.nuclphysa.2006.09.012).
- [15] C. Marquet, “Forward inclusive dijet production and azimuthal correlations in p(A) collisions,” *Nucl. Phys. A*, vol. 796, pp. 41–60, 2007, doi: [10.1016/j.nuclphysa.2007.09.001](https://doi.org/10.1016/j.nuclphysa.2007.09.001).

Backup

Wave Functions

- Momentum-space relation to the vacuum wave function:

$$\begin{aligned}\phi_0^\lambda(\mathbf{k}) &= 2\sqrt{1-y}[\delta_{\lambda,s} + (1-y)\delta_{\lambda,-s}] \frac{\mathbf{k} \cdot \boldsymbol{\varepsilon}_\lambda^*}{k^2} \\ \phi_-^\lambda(\mathbf{k}) &= \frac{k^2}{k^2 - M^2 - i\varepsilon} \phi_0^\lambda(\mathbf{k}) \\ \phi_+^\lambda(\mathbf{k}) &= \frac{k^2}{k^2 - M^2 + i\varepsilon} \phi_0^\lambda(\mathbf{k})\end{aligned}\tag{53}$$

- Coordinate-space

$$\phi_-^\lambda(\mathbf{u}) = \phi_Y^\lambda(\mathbf{u}) - \frac{1}{2}\phi_J^\lambda(\mathbf{u}), \phi_+^\lambda(\mathbf{u}) = \phi_Y^\lambda(\mathbf{u}) + \frac{1}{2}\phi_J^\lambda(\mathbf{u})\tag{54}$$

Wave Functions

$\phi_Y^\lambda \rightarrow \phi_0^\lambda$ as $M^2 \rightarrow 0$, while ϕ_J^λ is the on-shell discontinuity.